

PASCOLEUCYN Forte

Oral Drinkable Ampule

Indication:

Homeopathic remedy to support the immune system.

Ingredients (per 2 mL ampule):

Echinacea MT 18mg (supportive treatment in severe and fever infections), Aconitum napellus 4X 0.7mg (for highly acute inflammatory illnesses), Bryonia 4X 0.7mg (for acute inflammation of respiratory organs), Lachesis mutus 13X 0.7mg (for infectious diseases; inflammation of the skin and mucous membranes), Sulfur iodatum 4X 0.7mg (for inflammation of the airways and lungs; for lymph node/gland inflammation) in an isotonic sodium chloride solution.

Product Research Summary:

Pascoleucyn has a history of use dating back 60 years. It is considered **safe and effective for all ages** for the prevention and treatment of colds and flus. Studies have been conducted to establish its safety and efficacy in all age groups. Pascoleucyn is most effective when administered preventatively and also at the first signs of a cold or flu (headache, fever, sore throat, malaise, etc.)

In 1995 an observational study was conducted with 101 patients of medical and naturopathic doctors, aged 3-68 with either viral or bacterial respiratory infections. After 3 days, symptoms were reduced by 35%; after 7 days, symptoms were reduced by 80%.

Two retrospective studies took place in 2003. The first examined dosages for children, and involved 3,783 children aged 0-12 (including 235 infants) under the care of medical and naturopathic doctors. Therapists rated 100% effectiveness in all cases. Only 4 side effects were reported, consisting of a light rash.

Directions for Use:

Prevention – 1 ampule in Oct/Nov and 1 ampule in Jan/Feb or as directed by a naturopathic/medical doctor.
Onset of Flu – 1-2 ampules daily or as directed by a naturopathic/medical doctor.

Contraindications/Cautions:

Hypersensitivity to any of the active ingredients or excipients, or to members of the daisy family or to iodine. Pascoleucyn Forte must not be used in patients with diabetes mellitus, progressive systemic diseases such as tuberculosis, leukemia or leukemia-like conditions, inflammatory connective tissue disorders (collagen diseases), autoimmune disorders, multiple sclerosis, AIDS, HIV infection or any other chronic viral disease. Do not drink directly from vial due to sharp glass edges. In the absence of adequately documented experience, Pascoleucyn Forte should be used during pregnancy and breastfeeding only after consulting a health care practitioner. Do not use in children 0-12 years of age before a health care practitioner is consulted.

Interactions with other drugs:

None known.

Side Effects:

In isolated cases hypersensitive reactions may occur with drugs containing Echinacea. Skin rash, itching and in rare cases facial swelling, respiratory problems, dizziness and a fall in blood pressure have then been observed. In these cases please stop taking your medicine and consult with a health care practitioner.